

# Deep IR and optical studies of the fields of 3 proposed TŽO remnants

M.J.Coe and S.L.Pightling

*Department of Physics and Astronomy, The University, Southampton, SO17 1BJ, UK.*

Accepted . Received ;

## ABSTRACT

Deep infrared and optical images are presented of three proposed remnants of Thorne-Žytkow Objects (TŽO). In particular, the IR data go several infrared magnitudes deeper than previous observations and in at least one case reveal the existence of weak objects within the error circles. It is argued, however, that none of the objects is likely to be the binary companion to the X-ray source in that region. These data present severe limits on any possible star or residual envelope at the distances of the respective X-ray objects.

**Key words:** circumstellar matter – infrared: stars.

## 1 INTRODUCTION

Recent work by several authors, especially van Paradijs *et al.* (1995) and Mereghetti & Stella (1995), have identified a group of pulsating X-ray sources that appear to have no detectable optical counterpart. In addition, these objects show no evidence of any binarity and all have pulse periods in the 5–10s range. Their X-ray luminosities are all sufficiently high that accretion must be occurring on to the neutron star from some source other than the interstellar medium. One possible scenario is that they are the first objects identifiable as the remains of Thorne-Žytkow Objects (TŽOs).

The progenitor TŽOs would have consisted of a neutron star embedded in the core of another star (Thorne & Žytkow 1977). It is thought that these TŽOs could be formed as a result of the evolution of high mass systems through a common envelope phase (Ghosh *et al.* 1997). Once formed, thermonuclear burning via the  $r$ - $p$  process occurs in the core region of the TŽO. This supports the highly convective outer envelope against gravitational forces. However, once there is insufficient fuel remaining for the  $r$ - $p$  process to occur efficiently, the outer envelope collapses on the Kelvin-Helmholtz timescale (Podsiadlowski, Cannon and Rees, 1995). Some of the envelope material then forms into a disk around the neutron star with a radius that is dependent on the amount of angular momentum in the system. It is from this disk of material that the neutron star may accrete enough matter to power the X-ray pulsations seen.

In order to explore this possibility deep infrared imaging observations have been carried out from UKIRT and the NASA IRTF to search for any IR emission from such a disk around three of the neutron stars. The observations reach as deep as  $J \sim 20$  and in some cases reveal the presence of faint

objects in the X-ray error circles. However, the characteristics of the IR colours, and other considerations, strongly suggest that these are not the TŽO envelopes around the neutron stars. However, the new stringent limits on any IR companion in these systems leave these objects as enigmatic as ever.

The three systems investigated here are:

### 1.1 1E2259+586

The X-ray pulsar 1E2259+586 lies in the centre of the supernova remnant G109.1-1.0. The most recent estimates on its age make it approximately 3000 years old (Parmar *et al.* 1997), though values up to 10000 years have been quoted. G109.1-1.0 is at an estimated distance of about 4 kpc. Fahlman *et al.* (1982) demonstrated that the X-ray column densities to both the pulsar and the SNR were similar and hence that these two objects were probably related. However, the latest results from Parmar *et al.* (1997) indicate that this may not be the case. They find the X-ray column density to the pulsar to be greater than that to the supernova remnant. They suggest that this could be due to the pulsar being at a greater distance than the supernova remnant or to the presence of absorbing material local to the pulsar.

However, assuming the same distance to 1E2259+586 as the SNR implies a pulsar X-ray luminosity of  $\approx 10^{28}$  Watts. Results from *EXOSAT*, *GINGA* and *BeppoSAX* (Hanson *et al.* 1988, Koyama *et al.* 1989 and Parmar *et al.* 1997 respectively) have shown that the 7s period of 1E2259+586 is slowly increasing, but the spindown is too slow to power the observed luminosity. There is no evidence of binary motion,


the upper limit on  $a_x \sin i$  from pulse timing measurements being as low as 30 light-ms (Mereghetti, Israel and Stella, 1998). Early searches for the optical counterpart (Fahlman *et al.* 1982; Margon & Anderson 1983) found several possible candidates, one of which (star D) was tentatively identified with 1E2259+586 by Middleditch, Pennypacker & Burns (1983) on the basis of IR pulsations. However, more sensitive fast photometric observations by Davies *et al.* (1989) failed to find any pulsations, and there would now appear to be little positive evidence associating star D with 1E2259+586. A fainter, multicolour photometric study of all the possible counterparts by Davies & Coe (1991), together with revised astrometry, updated the list of possible counterparts.

Coe & Jones (1992) presented the first optical spectroscopy of all the candidates brighter than  $V=23$  and re-analysed the Einstein X-ray images of this source taken in 1981 in order to check the position and size of the X-ray error circle. They found that the relocation and reduction in the size of the error circle stemming from the re-processing of these data considerably changes the perspective on possible counterparts to the X-ray pulsar. In addition, a careful study of the images reveals previously unreported features that are possibly associated with jet activity from the pulsar, a possibility originally suggested by Gregory and Fahlman (1980). However, Hurford & Fesen 1995 report subsequent *Rosat* HRI observations which, while confirming the existence of such features, they suggest are just statistical fluctuations in the SNR emission.

## 1.2 4U0142+62

This persistent source has appeared in most X-ray catalogues and was originally confused (due to poor angular resolution) with the *Rosat* source RX J0146.9+6121 (=LSI +61 235) - the two sources are only 24 arcminutes apart. Measurements by White *et al.* (1987) with *EXOSAT* reported a 25m modulation from the region (later shown to be coming from LSI +61 235 (Mereghetti, Stella & De Nile 1993)) but using the unmodulated lower energy data they were able to identify the position of 4U0142+62. The detection of X-ray pulsations at 8.7s was reported by Israel *et al.* (1994) who concluded that they were most likely coming from the optically unidentified system 4U0142+62 rather than the new *Rosat* source. Further measurements by White *et al.* (1996) using *ASCA* confirmed this conclusion and also demonstrated evidence for an excess in the X-ray halo around the source. This could be due to material local to the pulsar along the line of site, possibly in the form of a molecular cloud or perhaps the remains of a TZO envelope. No evidence of any binary motion has been reported from the X-ray signal.

The accurate position reported by White *et al.* (1987) allowed them to locate the X-ray error circles from both *EXOSAT* and *Einstein* on the sky. As in the case of 1E2259+586 there is no evidence of any obvious optical candidates within the defined regions. They obtained an R band image and set a limit of  $R \geq 22.5$  for any counterpart. The *ASCA* observations (White *et al.* 1996) determined the column to the source to be  $8 \times 10^{21} \text{ cm}^{-2}$  which in turn allows them to deduce an optical extinction of  $A_v \sim 4.7$ . The distance to the source is not well defined but probably lies in the range 0.5–2.0 kpc. As in the case of 1E2259+586 a molecular cloud



**Figure 1.** A K band image of the field of 1E2259+586. The field size is 73 arcseconds and the circle shown represents the *Rosat* X-ray error circle of radius 5 arcsec (Dennerl, private communication). The three very faint objects within this error circle are referred to in the text as objects L, K and N from East to West respectively.

lies near to, or in front of 4U0142+62 (Mereghetti & Stella 1995) and may be affecting the column measurement.

## 1.3 RX J1838.4–0301

This source was originally reported by Schwentker (1994) as 5.5s pulsar possibly associated with a SNR at a distance of 4kpc. Interestingly, in this case the X-ray positional error circle was reported to include a  $\sim 14$  mag optical candidate. Subsequently Mereghetti *et al.* (1997) reanalysed the same *Rosat* data and reported a different interpretation of the data. They obtained an optical spectrum of the star in the error circle and concluded that this object was simply a K5 main sequence star and that the X-ray emission originated from its corona.

## 2 THE OBSERVATIONS

### 2.1 1E2259+586

This source was observed by UKIRT using IRCAM3 on 1996 October 23 in Service mode. The plate scale was  $\sim 0.3$  arcsec per pixel giving an image size of  $73 \times 73$  arcsec. Observations were taken in both J and K bands using a standard mosaicing technique with a total exposure of 27 minutes in each band. The K band image of the field is shown in Figure 1.

Superimposed on the image is the *Rosat* error circle (Dennerl, private communication) and the labels for several of the brighter stars in the region (following the convention

**Table 1.** Infrared observational measurements of the candidates for 1E2259+586

| Object | J band         | K band          | (J-K)           |
|--------|----------------|-----------------|-----------------|
| X      | $17.0 \pm 0.1$ | $16.1 \pm 0.06$ | $0.89 \pm 0.12$ |
| W      | $18.1 \pm 0.1$ | $17.0 \pm 0.1$  | $1.1 \pm 0.14$  |
| L      | $20.1 \pm 0.5$ | $17.9 \pm 0.4$  | $2.2 \pm 0.6$   |
| K      | $20.0 \pm 0.5$ | $18.0 \pm 0.4$  | $2.0 \pm 0.6$   |
| N      | $\geq 20.6$    | $18.3 \pm 0.4$  | $\geq 2.3$      |

of Coe et al. 1994a). The image is approximately 1-2 magnitudes deeper than any previous measurements and for the first time permits measurements of the three faint objects L, K and N.

In order to obtain an accurate calibration of the images, the established objects R1, R2, G, F, A and D (all shown in the figure) were used. The instrument magnitudes of these objects were compared with the reported fluxes in Davies & Coe (1991) and an excellent correlation was obtained (this, incidentally, confirms that there has been no detectable variation in the fluxes from any of these objects). Using this correlation the magnitudes for the three candidates L, K and N, as well as the nearby objects X and W, were determined. The results are presented in Table 1. From sampling the background around the area of interest the three sigma upper limits for any undetected source in this field may be determined to be  $J \geq 19.6$  and  $K \geq 18.4$ .


## 2.2 4U0142+62

This source was observed by the NASA IRTF on 1996 August 25 using the NSFCAM imager. This instrument has a plate scale of 0.3 arcsec per pixel giving an image size of 77 arcsec. Observations were taken in the J and K bands of 30 minutes each. In addition to the target field the standard star HD18881 was observed together with an appropriate number of flats.

The K band image of the field is shown in Figure 2. Also shown in this figure is the *Rosat* positional error circle from Hellier (1994). As one can see the field is extremely empty lacking any possible candidates close to the X-ray position. The faintest object detected and measured had a magnitude of  $J = 19.6 \pm 0.1$  and  $K = 16.88 \pm 0.07$ . The measurements of all the objects identified in the figure are given in Table 2 (note that star identification scheme is an extension of that started by White et al 1987).

## 2.3 RX J1838.4-0301

Notwithstanding the discussion in Section 1 of this paper, this source lacks a securely identified counterpart. Even the question of the existence, or otherwise, of X-ray pulsations needs resolving. It was therefore felt appropriate to explore the region around the error circle to see if any sources revealed themselves as typical of a counterpart to an X-ray source. Consequently, on 1996 November 15 images were taken in each of the B,V,R and I band using the Tek8 CCD camera on the 1m SAAO telescope with standard Johnson filters. In addition an image was recorded through the  $H\alpha$  filter. The exposure times for B,V and I were 300s, while R was 200s and  $H\alpha$  was 900s.




**Figure 2.** The upper panel shows a K band image of the field of 4U0142+62. The field size illustrated is 71 arcseconds and the circle shown represents the *Rosat* X-ray error circle of radius 3.2 arcsec (White et al. 1987). The lower panel is a V band image (Steinle, private communication) indicating the identification of stars in the region of the X-ray error circle.

If the X-ray pulsations are real then one obvious explanation for this system is that it is either a Be/X-ray binary system or a supergiant system. If this is the case then the optical counterpart would be a strong source of  $H\alpha$  emission. To investigate this possibility all the objects on the frame were measured in both the R band and  $H\alpha$  and the two sets of instrument magnitudes plotted against each other. This approach has been very successful in identifying counterparts to high mass X-ray binaries (see, for example, Coe et al., 1994b). In this case the result revealed no obvious source of  $H\alpha$  emission in the field.

**Table 2.** Infrared observational measurements of the candidates for 4U0142+62

| Object | J band       | K band       | (J-K)       |
|--------|--------------|--------------|-------------|
| 1      | 13.501±0.005 | 13.126±0.004 | 0.375±0.006 |
| 2      | 15.33±0.01   | 14.91 ±0.01  | 0.418±0.015 |
| 2a     | 17.61±0.04   | 17.09 ±0.03  | 0.513±0.05  |
| 3      | 15.92 ±0.02  | 15.16 ±0.01  | 0.76 ±0.02  |
| 4      | 15.14±0.01   | 14.86 ±0.01  | 0.27 ±0.01  |
| 6      | 14.95±0.01   | 14.63 ±0.01  | 0.32 ±0.01  |
| 8      | 15.74±0.01   | 15.44 ±0.01  | 0.30 ±0.02  |
| 9      | 15.52±0.01   | 15.20 ±0.01  | 0.32 ±0.02  |
| 13     | 16.82±0.02   | 16.38 ±0.02  | 0.43 ±0.03  |
| 14     | 17.15±0.03   | 16.67 ±0.03  | 0.47 ±0.04  |
| 16     | 17.02±0.03   | 16.38 ±0.02  | 0.64 ±0.03  |
| 25     | 18.55±0.06   | 17.54 ±0.04  | 1.01 ±0.07  |
| 29     | 17.05±0.03   | 16.65 ±0.02  | 0.40 ±0.04  |
| 30     | 17.16±0.03   | 16.34 ±0.02  | 0.82 ±0.03  |

**Figure 3.** K band image of RX J1838.4-0301. The field size is 73 arcseconds and the small circle of radius 5'' shows the estimate of the location of the *Rosat* source (Mereghetti et al., private communication and 1997), whereas the larger circle (radius 10'') is that of Schwentker (1994).

In addition to the above, deep IR observations were obtained from UKIRT using IRCAM3 on 1997 September 13 (J band) and 1997 September 29 (K band). Total exposures of 18 minutes were used in both bands. The instrument set up was identical to the observations of 1E2259+586. The K band image is shown in Figure 3 along with the determinations of the position of the *Rosat* source by Schwentker (1994) and Mereghetti et al. (private communication and 1997). The J and K magnitudes were measured of all the identified objects shown in Figure 3 and the results are presented in Table 3.

It is obvious from the UKIRT image that the reported "14th magnitude star" is in fact a complex of 5 or more previously unresolved stars. The limits on any undetected

**Table 3.** Infrared observational measurements of the candidates for RX J1838.4-0301

| Object | J band     | K band     | (J-K)      |
|--------|------------|------------|------------|
| A      | 12.69±0.03 | 12.74±0.03 | -0.05±0.04 |
| B      | 12.29±0.03 | 12.62±0.03 | -0.33±0.04 |
| C      | 14.01±0.03 | 14.69±0.03 | -0.68±0.04 |
| D      | 12.03±0.03 | 11.98±0.03 | +0.05±0.04 |
| E      | 13.72±0.03 | 14.16±0.03 | -0.44±0.04 |
| F      | 13.38±0.03 | 13.63±0.03 | -0.25±0.04 |
| G      | 13.36±0.03 | 13.24±0.03 | +0.12±0.04 |
| H      | 12.17±0.03 | 12.24±0.03 | -0.07±0.04 |
| J      | 11.55±0.03 | 12.01±0.03 | -0.46±0.04 |
| K      | 12.47±0.03 | 12.19±0.03 | +0.28±0.04 |
| L      | 13.19±0.03 | 14.15±0.03 | -0.96±0.04 |
| M      | 13.61±0.03 | 14.24±0.03 | -0.63±0.04 |

sources in the field were determined by sampling the background around the X-ray error circle. The resulting three sigma upper limits are  $J \geq 17.7$  and  $K \geq 16.9$ .

### 3 DISCUSSION

In order to investigate the nature of the objects in the fields of our targets an infrared colour plot was produced for each region. Figure 4 shows the plot for the first source, 1E2259+586.


To help interpret this diagram the location of the Main Sequence (MS) has been determined with the appropriate reddening. The solid line in the figure shows the MS with a reddening of  $A_v = 2.5$  (corresponding to a distance of 4kpc), while the dotted line is for  $A_v = 4.7$  (corresponding to 10kpc). These two values for reddening encompass the range of distances proposed for 1E2259+586, though the smaller value is the more probable (Davies & Coe 1991). As one can see, the reddening approximately follows the direction of the MS and hence does not provide a very useful distance discriminator. However, it is clear that most of the more obvious sources lie comfortably along the MS line and present no unusual IR colours. Object E, discussed in Coe & Jones (1992) as a possible white dwarf, is not detected in either IR image and hence is fainter than  $J=21$  and  $K=18.5$ .

The same cannot be said though for the new objects labelled K, L and N - these are the objects which actually lie in the *Rosat* error circle. A very large amount of reddening is required to make these objects consistent with a MS object - for example, a cluster of young B0 stars would need  $A_v \sim 13-14$ . Though it is highly improbable that the reddening in this galactic direction could, in general, be larger than  $A_v \sim 5$ , such a high value could be consistent with a large amount of local material around a distant cluster.

Alternatively,  $A_v = 4.7$  corresponds to  $E(J-K) = 0.8$  (Rieke & Lebofsky 1985) and the observed colour would be consistent with a Seyfert galaxy which typically has  $(J-K)_0 = 1.6$ , and hence would be reddened to  $(J-K) = 2.4$ . However, the chance of three such objects lying within 10 arcsec seems unlikely.

For the field containing 4U0142+61 the brightest 14 objects have had both their J and K colours determined. The results are presented in Figure 5.

Various fits to the data were undertaken to obtain the best estimate of the distance and reddening to the field




**Figure 4.** The infrared colours (J-K) for all the objects close to, or in, the X-ray error circle of 1E2259+586. The solid and dotted lines represent the location of the Main Sequence at reddening of  $A_v=2.5$  and  $4.7$  respectively.

stars. The solid line in the figure shows the best estimate for the Main Sequence at the distance (1kpc) and reddening ( $A_v=4.7$ ) quoted in White et al (1996). Clearly the field stars do not lie anywhere close to this position, nor does simply reducing the reddening to, for example,  $A_v=2$  (analogous perhaps to a position just in front of a molecular cloud reported by Mereghetti & Stella 1995). The best fit was obtained for an  $A_v=2.4$  at distance of 6kpc - indicated in Figure 5 by the dashed line. From this one can conclude that none of the visible stars are related to the X-ray source - perhaps not too surprising since they all lie well away from the X-ray error circle.

In the case of RX J1838.4-0301 the lack of any obvious  $H\alpha$  source in the immediate region of the X-ray position could be interpreted as support for the conclusions of Mereghetti et al. (1997) that the X-ray signal is coronal in nature. However, the IR images reported here clearly show the presence of several comparably bright stars in the region of the *Rosat* source. Figure 6 shows a similar plot to Figures 4 and 5, but for the objects in the field of RX J1838.4-0301.


Schwentker (1994) reports a hydrogen column to this source in the range  $(1-3) \times 10^{21}$  atoms per sq.cm. If we use this value with the relationship given in Ryter, Cesarsky & Audouze (1975) we can deduce an  $E(B-V)$  in the range 0.2 -



**Figure 5.** The infrared colours (J-K) for all the objects close to the X-ray error circle of 4U0142+62. The solid line represent the location of the Main Sequence at a reddening of  $A_v=4.7$  and a distance of 1kpc, and the dotted line the MS with  $A_v=2.4$  at distance of 6kpc.

0.4 and hence an  $E(J-K)$  in the range 0.1 - 0.2. The locus of the Main Sequence at an average reddening of  $E(J-K)=0.15$  (i.e.  $E(B-V)=0.27$ ) and the distance of 1kpc suggested by Schwentker (1994) is marked by the dashed line in Figure 6. Many of the stars in the region of the X-ray source lie close to this line, but the spread is quite significant in the (J-K) axes. Mereghetti et al (1997) have proposed a K5 star at 300pc as being consistent with the X-ray data (if the pulsations are not real). A study of all the stars in this direction indicates that at distances beyond 200-300 pc the hydrogen column will always be of the order  $10^{21}$  atoms per sq.cm. The solid line in Figure 6 shows the reddening expected for such a column, but now at a distance of 300pc. Star A lies extremely close to the position expected for a K5 star at such a distance and reddening, and this object may have been the dominating star in Mereghetti et al's data. However, the other stars in the *Rosat* error circle lie scattered all over the diagram and offer many other possible solutions. For example, Star D is the brightest in our IR data at  $J=12.0$  and probably lies even closer than Star A. Certainly Mereghetti et al might well have had several of the stars A-E in the 2" slit for their spectroscopy.

If the pulsations turn out to be real as proposed by Schwentker (1994), then it would support RX J1838.4-0301



**Figure 6.** The infrared colours ( $J-K$ ) for all the objects close to, or in, the X-ray error circle of RX J1838.4-0301. The solid line represent the location of the Main Sequence at a reddening of  $A_v=0.42$  and a distance of 0.3kpc. The dashed line is the MS at the same  $A_v$ , but at a distance of 1.0kpc.

remaining a member of this class of enigmatic objects. What is clear is that it is very unlikely to be a member of any Be or supergiant binary system. Further X-ray observations are clearly required both to refine the X-ray position in this crowded field, and to clarify the situation regarding the possible X-ray pulsations.

Finally, a rough estimate of the limiting size for a disk producing black body emission may be obtained from the IR results presented here. In the case of 1E2259+586 the limits of  $J \geq 20$  and  $K \geq 18.5$  imply a limiting size of approximately 1 solar radius ( $7 \times 10^8$  m) for a 2500K black body at 4.5kpc. Podsiadlowski, Cannon and Rees (1995) quote a size of the order  $3 \times 10^8$  m for a system that had an initial binary orbital period of 10d and a companion of 15 solar masses. The IR limits for 4U0142+62 are similar and imply a somewhat smaller disk limit if the distance to the source is at the lower end of the proposed range of (i.e. 0.5 kpc) - maybe by up to a factor of 10 (i.e. a disk size of  $7 \times 10^7$  m). However, the upper bounds to the distance (1.5 kpc) give limits similar to the disk sizes of Podsiadlowski, Cannon and Rees (1995). Thus these new IR values are probing the expected disk sizes for a proposed TZO envelope.

## 4 CONCLUSIONS

The first deep infrared and optical images have been presented of three proposed remnants of Thorne-Żytkow Objects (TZO). In particular, the IR data go several infrared magnitudes deeper than previous observations and in at least one case, 1E2259+586, reveal the existence of weak objects within the X-ray error circle. It is argued, however, that in that case none of the objects seen is likely to be the binary companion to the X-ray source in that region. These data present new limits on any possible IR counterpart to the respective X-ray objects.

## ACKNOWLEDGMENTS

We are very grateful to UKIRT for obtaining data on two of these sources through their Service programme, and also to the NASA IRTF for the data on 4U0142+62 obtained in a similar manner. We thank Dave Buckley for obtaining the optical CCD frames of RX J1838.4-0301 and to Sandro Mereghetti for many helpful comments. SLP acknowledges the support of a PPARC studentship.

## REFERENCES

- Coe M.J., Jones, L.R. 1992, MNRAS, 259, 191.
- Coe M.J., Jones L.R., Lehto H., 1994a, MNRAS 270, 178.
- Coe M.J., Roche P., Everall C. et al., 1994b, MNRAS 270, L57.
- Davies S.R., Coe M.J., Payne B.J., Hanson C.G., 1989, MNRAS, 237, 973
- Davies S.R., Coe M.J., 1991, MNRAS, 249, 313
- Fahlman G.G., Hickson P., Richer H.B., Gregory P.C., Middleditch J., 1982, ApJ, 261, L1
- Gregory P.C., Fahlman G.G., 1980 Nat, 287, 805
- Hanson C.G., Dennerl K., Coe M.J., Davies S.R., 1988, A&A, 195, 114
- Hellier C., 1994, MNRAS 271, L21.
- Hurford A.P. & Fesen R.A., 1995 MNRAS 277, 549.
- Israel G.L., Mereghetti S., Stella L., 1994, ApJ 433, L25.
- Koyama K. et al., 1989, PASJ, 41, 461
- Margon B., Anderson S.F., 1983, Astrophys. Lett., 23, 211
- Mereghetti S., Stella L., De Nile F., 1993 A&A 278, L23.
- Mereghetti S., Stella L., 1995, ApJ 442, L17.
- Mereghetti S., Belloni T., Nasuti F.P., 1997, A&A 321, 835.
- Mereghetti S., Stella L., Israel G.L., 1998 MNRAS In press.
- Middleditch J., Pennypacker C.R., Burns, M.S. 1983, ApJ, 274, 313
- Parmar A.N., Oosterbroek T., Favata F., 1997, A&A submitted.
- Podsiadlowski P., Cannon R.C., Rees M.J. 1995 MNRAS 274, 485.
- Rieke G.H., Lebofsky M.J., 1985, ApJ 288, 618.
- Ryter C., Cesarsky C.J., Audouze J. 1975, ApJ 198, 103.
- Schwentker O., 1994, A&A 286, L47.
- Thorne K.S., Zytkow A.N., 1977, ApJ 212, 832.
- van Paradijs J., Taam R.E., van den Heuvel E.P.J., 1995, A&A 299, L41.
- White N.E., Mason K.O., Giommi P., Angelini L., Pooley G., Murdin P.G., Brabduardi-Raymont G., Wall J.V. 1987 MNRAS 226, 645.
- White N.E., Angelini L., Ebisawa K., Tanaka Y., Ghosh P. 1996 ApJ 463, L83.

This paper has been produced using the Royal Astronomical  
Society/Blackwell Science L<sup>A</sup>T<sub>E</sub>X style file.